arXiv:1503.05397v1 [math.CA] 26 Jan 2015

# ADDENDUM TO : ORTHONORMAL BASES OF REGULAR WAVELETS IN SPACES OF HOMOGENEOUS TYPE

PASCAL AUSCHER AND TUOMAS HYTÖNEN*

Abstract. We bring a precision to our cited work concerning the notion of "Borel measures", as the choice among different existing definitions impacts on the validity of the results.

We wish to bring a precision to our work [1]. The same remarks apply to the follow-up article [2]. Proposition 4.5 in [1] states that if $\mu$ is a non-trivial Borel measure on a quasi-metric space $X$, finite on bounded Borel sets, and $1 \leq p < \infty$, then Hölder-$\eta$-continuous functions of bounded support are dense in $L^p(\mu)$, where $\eta$ is the Hölder exponent of the splines constructed in [1].

Depending on the meaning of "Borel measure", as different definitions can be found in the literature, the result is correct or wrong.

If $\mu$ is a $\sigma$-additive measure on the Borel $\sigma$-algebra, then the result with the given proof is correct, as the measurable sets coincide with the Borel sets. In that case, all our results are valid as stated.

However, if $\mu$ is an outer measure or a $\sigma$-additive measure on $X$ for which the Borel sets are $\mu$-measurable, then for the first sentence of the proof to be valid one must add the condition that for every $\mu$-measurable set $A$ (in the sense of Caratheodory for the outer measure case), there is a Borel set $B \supseteq A$ such that $\mu(A) = \mu(B)$, and the rest of the proof goes through. In [3] for example, Borel outer measures are called regular if this condition holds for all $A$ (not necessarily $\mu$-measurable). With such a definition of a Borel measure, this regularity condition should be added to our statements. Without regularity, the correct conclusion of Proposition 4.5 is density in the space of $L^p$ functions having a Borel measurable representative. Thus, our wavelet representations are valid for functions in this subspace and $1 < p < \infty$. This is enough for many purposes.

## References

[1] Auscher, P. and Hytönen, T. Orthonormal bases of regular wavelets in spaces of homogeneous type, *Appl. Comput. Harmon. Anal.*, 34 (2013), 266–296.

[2] Hytönen, T. and Tapiola, O. Almost Lipschitz-continuous wavelets in metric spaces via a new randomization of dyadic cubes, *J. Approx. Theory*, 185 (2014), 12–30.

[3] Mattila, P. Geometry of sets and measures in euclidean spaces". Cambridge Studies in Advanced Mathematics, 44. Cambridge University Press, Cambridge, 1995.

Laboratoire de Mathématiques, UMR 8628, Univ. Paris-Sud and CNRS, F-91405 Orsay
*E-mail address*: `pascal.auscher@math.u-psud.fr`

Department of Mathematics and Statistics, P.O. Box 68, FI-00014 University of Helsinki, Finland
*E-mail address*: `tuomas.hytonen@helsinki.fi`

---

2010 *Mathematics Subject Classification.* 42C40, 41A15, 30Lxx, 42B25.
*Key words and phrases.* Quasi-metric space, Borel measure, spline, wavelet.
*Corresponding author. Telephone: +358 2941 51430.